arXiv:1101.2611v1 [math.NT] 13 Jan 2011

IDENTITIES INVOLVING VALUES OF BERNSTEIN, q-BERNOULLI AND q-EULER POLYNOMIALS

A. BAYAD AND T. KIM

Abstract In this paper we give some relations involving values of q-Bernoulli, q-Euler and Bernstein polynomials. From these relations, we obtain some interesting identities on the q-Bernoulli, q-Euler and Bernstein polynomials.

1. Introduction and preliminaries

Throughout this paper, let p be a fixed odd prime number. The symbols, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ denote the ring of p-adic integers, the field of p-adic numbers and the field of p-adic completion of the algebraic closure of $\mathbb{Q}_p$, respectively. When one talks of q-extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$, or a p-adic $q \in \mathbb{C}_p$. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let $x = p^r \frac{s}{t}$ where $r \in \mathbb{Q}$ and $(s,t) = (p,s) = (p,t) = 1$. Then the p-adic absolute value is defined by $|x|_p = p^{-r}$. If $q \in \mathbb{C}$, we assume $|q| < 1$, and if $q \in \mathbb{C}_p$, we always assume $|1-q|_p < 1$.

Let $C_{p^n} = \{\xi \; |\xi^{p^n} = 1\}$ be the cyclic group of order p^n. Then, the p-adic locally constant space is defined as $T_p = \lim_{n\to\infty} C_{p^n}$ (see [10]). Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the bosonic p-adic invariant integral on $\mathbb{Z}_p$ is defined by

$$\begin{aligned} I_1(f) = \int_{\mathbb{Z}_p} f(x)d\mu(x) &= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\mu(x+p^N\mathbb{Z}_p) \qquad (1) \\ &= \lim_{N\to\infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x), \quad \text{(see [17, 10, 11, 12])}. \end{aligned}$$

From (1), we note that

$$I_1(f_1) = \int_{\mathbb{Z}_p} f(x+1)d\mu(x) = \int_{\mathbb{Z}_p} f(x)d\mu(x) + f'(0), \qquad (2)$$

2000 Mathematics Subject Classification : 11B68, 11B73, 41A30.

Key words and phrases : Bernstein polynomial, q-bernoulli numbers and polynomials, q-Euler numbers and polynomials, fermionic p-adic intergal, bermionic p-adic intergal .

where $f_1(x) = f(x+1)$, (see [14, 10]).
Let $f \in UD(\mathbb{Z}_p)$. Then the fermionic integral on $\mathbb{Z}_p$ is given by

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\mu_{-1}(x+p^N\mathbb{Z}_p) \tag{3}$$
$$= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x, \quad \text{(see [18, 17, 9]).}$$

As known results, by (3), we get

$$I_{-1}(f_1) = -I_{-1}(f) + 2f(0), \text{ (see [15]).} \tag{4}$$

By using (2) and (4), we investigate some properties for the q-Bernoulli polynomials and the q-Euler polynomials.
Let $C[0,1]$ be denote by the space of continuous functions on $[0,1]$. For $f \in C[0,1]$, Bernstein introduced the following well-known linear positive operator in the field of real numbers $\mathbb{R}$:

$$\mathbb{B}_n(f|x) = \sum_{k=0}^{n} f(\frac{k}{n})\binom{n}{k}x^k(1-x)^{n-k} = \sum_{k=0}^{n} f(\frac{k}{n})B_{k,n}(x),$$

where $\binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{k!} = \frac{n!}{k!(n-k)!}$ (see [1, 3, 18, 17, 14, 12]). Here, $\mathbb{B}_n(f|x)$ is called the Bernstein operator of order n for f. For $k, n \in \mathbb{Z}_+$, the Bernstein polynomials of degree n are defined by

$$B_{k,n}(x) = \binom{n}{k}x^k(1-x)^{n-k}, \quad \text{for } x \in [0,1]. \tag{5}$$

For $x \in \mathbb{Z}_p$, the p-adic extension of Bernstein polynomials are given by

$$B_{k,n}(x) = \binom{n}{k}x^k(1-x)^{n-k}, \text{ where } k, n \in \mathbb{Z}_+, \text{ (see [18, 22, 5].} \tag{6}$$

The purpose of this paper is to give some relations for the q-Bernoulli, q-Euler and Bernstein polynomials.
From these relations, we obtain some interesteing identities on the q-Bernoulli, q-Euler and Bernstein polynomials.

2. SOME IDENTITIES ON THE q-BERNOULLI, q-EULER AND BERNSTEIN POLYNOMIALS

In this section we assume that $q \in T_p$. Let $f(x) = q^x e^{xt}$. From (1) and (2), we have

$$\int_{\mathbb{Z}_p} f(x) d\mu(x) = \frac{t}{qe^t - 1}, \text{ (see [10]).} \tag{7}$$

The q-Bernoulli numbers are defined by

$$\frac{t}{qe^t-1} = e^{B(q)t} = \sum_{n=0}^{\infty} B_n(q)\frac{t^n}{n!}, \text{ (see [10]),} \tag{8}$$

with usual convention about replacing $B_n(q)$ by $B_n(q)$.
By (7) and (8), we get Witt's formula for the q-Bernoulli numbers as follows:

$$\int_{\mathbb{Z}_p} q^x x^n d\mu(x) = B_n(q),\ n \in \mathbb{Z}_+,\ (\text{ see [10]}). \tag{9}$$

From (8), we can derive the following recurrence sequence:

$$B_0(q) = 0,\ \text{and}\ q(B(q)+1)^n - B_n(q) = \begin{cases} 1 & \text{if}\ \ n = 1, \\ 0 & \text{if}\ \ n > 1, \end{cases} \tag{10}$$

with usual convention about replacing $B^n(q)$ by $B_n(q)$.
Now, we define the q-Bernoulli polynomials as follows:

$$\frac{t}{qe^t - 1} e^{xt} = e^{B(x|q)t} = \sum_{n=0}^{\infty} B_n(x|q) \frac{t^n}{n!}, \tag{11}$$

with usual convention about replacing $B^n(x|q)$ by $B_n(x|q)$.
By (2) and (11), we easily get

$$\int_{\mathbb{Z}_p} q^y e^{(x+y)t} d\mu(y) = \frac{t}{qe^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x|q) \frac{t^n}{n!}. \tag{12}$$

Thus, we also obtain Witt's formula for the q-Bernoulli polynomials as follows:

$$\int_{\mathbb{Z}_p} q^y (x+y)^n d\mu(y) = B_n(x|q), \quad \text{for}\ n \in \mathbb{Z}_+. \tag{13}$$

By (9) and (13), we easly get

$$\begin{aligned} B_n(x|q) &= \sum_{l=0}^{n} \binom{n}{l} x^{n-l} B_l(q) \\ &= (x + B(q))^n, \quad \text{for}\ n \in \mathbb{Z}_+. \end{aligned}$$

It is easy to show that

$$\begin{aligned} \frac{t}{qe^t - 1} &= \left(\frac{t}{q(1-q^{-1})} \right) \left(\frac{1-q^{-1}}{e^t - q^{-1}} \right) \\ &= \frac{1}{q(1-q^{-1})} \sum_{n=0}^{\infty} H_n(q^{-1})(n+1) \frac{t^{n+1}}{(n+1)!}, \end{aligned} \tag{14}$$

where $H_n(q^{-1})$ are the n-th Frobenius-Euler numbers (see [17]).
By (8), (10) and (15), we get

$$\frac{B_{n+1}(q)}{n+1} = \frac{H_n(q^{-1})}{q(1-q^{-1})} = \frac{H_n(q^{-1})}{q-1}. \tag{15}$$

Therefore, by (15), we obtain the following proposition.

Proposition 1. *For any $n \in \mathbb{Z}_+$, we have*

$$\frac{B_{n+1}(q)}{n+1} = \frac{H_n(q^{-1})}{q(1-q^{-1})} = \frac{H_n(q^{-1})}{q-1},$$

where $H_n(q^{-1})$ are the n-th Frobenius-Euler numbers.

From (11) and (12), we can derive the following equation:

$$\frac{qt}{qe^t-1}e^{(1-x)t}=\frac{-t}{q^{-1}e^{-t}-1}e^{-xt}=\sum_{n=0}^{\infty}B_n(x|q^{-1})(-1)^n\frac{t^n}{n!}, \tag{16}$$

and

$$\frac{qt}{qe^t-1}e^{(1-x)t}=qe^{B(1-x|q)t}=q\sum_{n=0}^{\infty}B_n(1-x|q)\frac{t^n}{n!}. \tag{17}$$

By comparing the coefficients on the both sides of (16) and (17), we obtain the following theorem.

Theorem 2. *Let $n\in\mathbb{Z}_+$. Then we have*

$$qB_n(1-x|q)=(-1)^nB_n(x|q^{-1}).$$

From (10) and (13), we have

$$\begin{aligned} B_n(2|q) &= (B(q)+1+1)^n=\sum_{l=0}^{n}\binom{n}{l}B_l(1|q) \\ &= B_0(q)+\frac{1}{q}\sum_{l=1}^{n}\binom{n}{l}qB_l(1|q)=\frac{1}{q}\sum_{l=1}^{n}\binom{n}{l}B_l(q) \\ &= \frac{1}{q}B_n(1|q)=\frac{1}{q^2}qB_n(1|q)=\frac{1}{q^2}B_n(q), \text{ while } n>1. \end{aligned} \tag{18}$$

Therefore, by (18), we obtain the following theorem.

Theorem 3. *For $n\in\mathbb{Z}_+$ with $n>1$, we have*

$$q^2B_n(2|q)=B_n(q).$$

By (9) and (13), we easily see that

$$\begin{aligned} \int_{\mathbb{Z}_p}q^{-x}(1-x)^nd\mu(x) &= (-1)^n\int_{\mathbb{Z}_p}q^{-x}(x-1)^nd\mu(x) \\ &= q\int_{\mathbb{Z}_p}(x+2)^nq^xd\mu(x) \\ &= \frac{1}{q}\int_{\mathbb{Z}_p}x^nq^xd\mu(x). \end{aligned} \tag{19}$$

Therefore, by (19), we obtain the following theorem.

Theorem 4. *For $n\in\mathbb{N}$ with $n>1$, we have*

$$\int_{\mathbb{Z}_p}q^{-x}(1-x)^nd\mu(x)=\frac{1}{q}\int_{\mathbb{Z}_p}x^nq^xd\mu(x).$$

Let $f(x)=q^xe^{xt}$. Then, from (3) and (4), we note that

$$\int_{\mathbb{Z}_p}q^xe^{xt}d\mu_{-1}(x)=\frac{2}{qe^t+1},\ (\text{ see [18, 19, 4, 21]}). \tag{20}$$

In [18], the q-Euler numbers are defined by

$$\frac{2}{qe^t+1}=e^{E(q)t}=\sum_{n=0}^{\infty}E_n(q)\frac{t^n}{n!}, \tag{21}$$

with usual convention about replacing $E^n(q)$ by $E_n(q)$.
Let us define the q-Euler polynomials as follows:

$$\frac{2}{qe^t+1}e^{xt} = e^{E(x|q)t} = \sum_{n=0}^{\infty} E_n(x|q)\frac{t^n}{n!}. \tag{22}$$

From (3) and (4), we note that

$$\int_{\mathbb{Z}_p} q^y e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{qe^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x|q)\frac{t^n}{n!}. \tag{23}$$

Thus, by (22) and (23), we get

$$\int_{\mathbb{Z}_p} q^y (x+y)^n d\mu_{-1}(y) = E_n(x|q), \quad \text{for } n \in \mathbb{Z}_+. \tag{24}$$

By (20), (21) and (24), we get

$$E_n(x|q) = \sum_{l=0}^{n} \binom{n}{l} x^{n-l} E_{l,q} = (x+E(q))^n\,, \tag{25}$$

with usual convention $E(q)^n$ by $E_n(q)$.
In [2], it is known that

$$qE_n(2|q) = 2 + \frac{1}{q}E_n(q), \tag{26}$$

and

$$\int_{\mathbb{Z}_p} q^{-x}(1-x)^n d\mu_{-1}(x) = 2 + \frac{1}{q}\int_{\mathbb{Z}_p} x^n q^x d\mu_{-1}(x), \text{ (see [13]).}$$

For $x \in \mathbb{Z}_p$, by (6), we have

$$\begin{aligned} B_n(1-x|q) &= \sum_{l=0}^{n} \binom{n}{l} (1-x)^{n-l} B_l(q) \\ &= \sum_{l=0}^{n} \binom{n}{l} (1-x)^{n-l} x^l B_l(q) x^{-l} \\ &= \sum_{l=0}^{n} B_{l,n}(x) B_l(q) x^{-l}, \end{aligned} \tag{27}$$

where $B_{l,n}(x)$ are Bernstein polynomials of degree n.
Therefore, by (27), we obtain the following proposition.

Proposition 5. *For $n \in \mathbb{Z}_+$, we have*

$$B_n(1-x|q) = \sum_{l=0}^{n} B_{l,n}(x) B_l(q) x^{-l},$$

where $B_{l,n}(x)$ are Bernstein polynomials of degree n.
From (7) and (20), we can derive the following equation:

$$\int_{\mathbb{Z}_p}\int_{\mathbb{Z}_p} q^{x+y} e^{(x+y)t} d\mu_{-1}(x) d\mu(y) = \frac{2t}{q^2e^{2t}-1}. \tag{28}$$

By (8), we get

$$\frac{2t}{q^2e^{2t}-1}=e^{B(q^2)2t}=\sum_{n=0}^{\infty}2^nB_n(q^2)\frac{t^n}{n!}. \tag{29}$$

By comparing the coefficients on the both sides of (28) and (29), we obtain the following theorem.

Theorem 6. *For $n\in\mathbb{Z}_+$, we have*

$$\frac{1}{2^n}\int_{\mathbb{Z}_p}\int_{\mathbb{Z}_p}q^{x+y}(x+y)^{n+1}d\mu_{-1}(x)d\mu(y)=B_n(q^2).$$

By using (9) and (24), we obtain the following corollary.

Corollary 7. *For $n\in\mathbb{Z}_+$, we have*

$$\frac{1}{2^n}\sum_{l=0}^{n}\binom{n}{l}E_l(q)B_{n-l}(q)=B_n(q^2).$$

From the definition of Bernstein polynomials, we note that

$$B_{k,n}(x)=B_{n-k,n}(1-x),\quad \text{for } n,k\in\mathbb{Z}_+.$$

Thus, we have

$$B_{k,n}(\frac{1}{2})=B_{n-k,n}(\frac{1}{2})=\binom{n}{k}\left(\frac{1}{2}\right)^{n-k}\left(\frac{1}{2}\right)^{k}=\left(\frac{1}{2}\right)^{n}\binom{n}{k}.$$

Therefore, we obtain the following lemmma.

Lemma 8. *Let $n,k\in\mathbb{Z}_+$. Then we have*

$$B_{k,n}(\frac{1}{2})=\left(\frac{1}{2}\right)^{n}\binom{n}{k}.$$

By Lemma 8 and Corollary 7, we obtain the following corollary.

Corollary 9. *For $n\in\mathbb{Z}_+$, we have*

$$\begin{aligned}B_n(q^2)&=\frac{1}{2^n}\sum_{l=0}^{n}\binom{n}{l}E_l(q)B_{n-l}(q)\\&=\sum_{l=0}^{n}B_{l,n}(\frac{1}{2})E_l(q)B_{n-l}(q).\end{aligned}$$

For the right side of (28), we have

$$\begin{aligned}\frac{2t}{q^2e^{2t}-1}&=-2t\sum_{m=0}^{\infty}q^{2m}e^{2mt}=-2t\sum_{n=0}^{\infty}\left(2^n\sum_{m=0}^{\infty}q^{2m}m^n\right)\frac{t^n}{n!}\\&=\sum_{n=0}^{\infty}\left(-2^{n+1}(n+1)\sum_{m=0}^{\infty}q^{2m}m^n\right)\frac{t^{n+1}}{(n+1)!}.\end{aligned} \tag{30}$$

In (29), we see that $B_0(q^2)=0$. By comparing coefficients on the both sides of (29) and (30), we obtain the following theorem.

Theorem 10. *For $n \in \mathbb{Z}_+$, we have*

$$-\frac{1}{2^{(n+1)}(n+1)} \int_{\mathbb{Z}_p}\int_{\mathbb{Z}_p} q^{x+y}(x+y)^n d\mu_{-1}(x)d\mu(y) = \sum_{m=0}^{\infty} q^{2m}m^n.$$

By (9), (24) and binomial theorem, we obtain the following corollary

Corollary 11. *For $n \in \mathbb{Z}_+$, we have*

$$\sum_{m=0}^{\infty} q^{2m}m^n = -\frac{1}{2^{(n+1)}(n+1)} \sum_{l=0}^{n+1} \binom{n+1}{l} E_l(q)B_{n+1-l}(q).$$

By Lemma 7, we obtain the following corollary.

Corollary 12. *For $n \in \mathbb{Z}_+$, we have*

$$\sum_{m=0}^{\infty} q^{2m}m^n = -\frac{1}{n+1} \sum_{l=0}^{n+1} B_{n+1,l}(\frac{1}{2})E_l(q)B_{n+1-l}(q).$$

It seems to be important to study double p-adic integral representation of bosonic and fermionic on the Bernstein polynomials associated with q-Bernoulli and q-Euler polynomials. Theorem 10 is useful to study those integral representation on $\mathbb{Z}_p$ related to Bernstein polynomials. Now, we take bosonic p-adic invariant integral on $\mathbb{Z}_p$ for one Bernstein polynomials in (6)as follows:

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x)q^x d\mu(x) &= \int_{\mathbb{Z}_p} \binom{n}{k}(1-x)^{n-k}x^k q^x d\mu(x) \\ &= \binom{n}{k}\sum_{l=0}^{n-k}\binom{n-k}{l}(-1)^l \int_{\mathbb{Z}_p} x^{k+l}q^x d\mu(x) \\ &= \binom{n}{k}\sum_{l=0}^{n-k}\binom{n-k}{l}(-1)^l B_{k+l}(q). \end{aligned} \tag{31}$$

From the reflection symmetry of Bernstein polynomials, we have

$$B_{k,n}(x) = B_{n-k,n}(1-x), \quad \text{where } n,k \in \mathbb{Z}_+. \tag{32}$$

Let $n, k \in \mathbb{N}$ with $n > k+1$. Then, by (32), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x)q^x d\mu(x) &= \int_{\mathbb{Z}_p} B_{n-k,n}(1-x)q^x d\mu(x) \\ &= \binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k-l}\int_{\mathbb{Z}_p}(1-x)^{n-l}q^x d\mu(x) \\ &= q\binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k-l}\int_{\mathbb{Z}_p} q^{-x}x^{n-l}d\mu(x) \\ &= q\binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k-l}B_{n-l}(q^{-1}). \end{aligned} \tag{33}$$

Therefore, we obtain the following theorem.

Theorem 13. *Let $n, k \in \mathbb{Z}_+$ with $n > k+1$. Then we have*

$$\int_{\mathbb{Z}_p} B_{k,n}(x) q^{1-x} d\mu(x) = 2^k \binom{n}{k} \sum_{l=0}^{k} B_{l,k}(\frac{1}{2})(-1)^{k-l} B_{n-l}(q).$$

By (31), we obtain the following corollary

Corollary 14. *For $n, k \in \mathbb{Z}_+$ with $n > k+1$, we have*

$$2^{2k}\binom{n}{k} B_{l,k}(\frac{1}{2})(-1)^{k-l} B_{n-l}(q) = 2^n q \sum_{l=0}^{n-k} B_{l,n-k}(\frac{1}{2})(-1)^l B_{k+l}(q).$$

Let $m, n, k \in \mathbb{Z}_+$, with $m+n > k+1$. Then we have

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) q^{-x} d\mu(x) &= \binom{n}{k}\binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \int_{\mathbb{Z}_p} q^{-x}(1-x)^{n+m-l} d\mu(x) \\ &= \binom{n}{k}\binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} q \int_{\mathbb{Z}_p} q^{x}(x+2)^{n+m-l} d\mu(x) \\ &= \binom{n}{k}\binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \frac{1}{q} B_{n+m-l}(q). \end{aligned} \tag{34}$$

Therefore, by (34), we obtain the following theorem.

Theorem 15. *Let $m, n, k \in \mathbb{Z}_+$ with $m+n > k+1$. Then we have*

$$\int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) q^{1-x} d\mu(x) = \binom{n}{k}\binom{m}{k} 2^{2k} \sum_{l=0}^{2k} B_{l,2k}(\frac{1}{2})(-1)^{l+2k} B_{n+m-l}(q).$$

By binomial theorem, we easily get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) q^{1-x} d\mu(x) &= \binom{n}{k}\binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \int_{\mathbb{Z}_p} x^{l+2k} q^{1-x} d\mu(x)) \\ &= q\binom{n}{k}\binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l B_{l+2k}(\frac{1}{q}). \end{aligned} \tag{35}$$

By (35), we obtain the following corollary.

Corollary 16. *Let $m, n, k \in \mathbb{Z}_+$ with $n+m > 2k+1$. Then we have*

$$2^{4k} \sum_{l=0}^{2k} (-1)^{l+2k} B_{l,2k}(\frac{1}{2}) B_{n+m-l}(q) = 2^{n+m} q \sum_{l=0}^{n+m-2k} (-1)^l B_{l,n+m-2k}(\frac{1}{2}) B_{l+2k}(\frac{1}{q}).$$

For $s \in \mathbb{N}$, let $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$ with $n_1 + n_2 + \cdots + n_s > sk + 1$.

$$\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n_1}(x) \cdots B_{k,n_s}(x) q^{-x} d\mu(x) = \binom{n_1}{k} \cdots \binom{n_s}{k} \int_{\mathbb{Z}_p} x^{sk}(1-x)^{n_1+\ldots+n_s} q^{-x} d\mu(x) \\
= \quad & \binom{n_1}{k} \cdots \binom{n_s}{k} \sum_{l=0}^{sk} (-1)^{sk+l} \int_{\mathbb{Z}_p} (1-x)^{n_1+\ldots+n_s-l} q^{-x} d\mu(x) \\
= \quad & q\binom{n_1}{k} \cdots \binom{n_s}{k} \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \int_{\mathbb{Z}_p} (x+2)^{n_1+\ldots+n_s-l} q^{x} d\mu(x) \\
= \quad & \frac{1}{q}\binom{n_1}{k} \cdots \binom{n_s}{k} \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} B_{n_1+\cdots+n_s-l}(q). \qquad (36)
\end{aligned}$$

Therefore, by (36), we obtain the following theorem.

Theorem 17. *For $s \in \mathbb{N}$, let $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$ with $n_1+n_2+\cdots+n_s > sk+1$. Then we have*

$$\int_{\mathbb{Z}_p} B_{k,n_1}(x) \cdots B_{k,n_s}(x) q^{1-x} d\mu(x) = \binom{n_1}{k} \cdots \binom{n_s}{k} 2^{sk} \sum_{l=0}^{sk} (-1)^{sk+l} B_{l,sk}(\frac{1}{2}) B_{n_1+\cdots+n_s-l}(q).$$

From binomial theorem, we can easily derive the following equation:

$$\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n_1}(x) \cdots B_{k,n_s}(x) q^{-x} d\mu(x) \qquad (37) \\
&= \binom{n_1}{k} \cdots \binom{n_s}{k} \sum_{l=0}^{n_1+\cdots+n_s-sk} \binom{n_1+\cdots+n_s-sk}{l} (-1)^l \int_{\mathbb{Z}_p} x^{l+sk} q^{-x} d\mu(x) \\
&= \binom{n_1}{k} \cdots \binom{n_s}{k} \sum_{l=0}^{n_1+\cdots+n_s-sk} \binom{n_1+\cdots+n_s-sk}{l} (-1)^l B_{l+sk}(q^{-1}).
\end{aligned}$$

By (37), we obtain the following corollary.

Corollary 18. *Let $s \in \mathbb{N}$ and $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$ with $n_1+n_2+\cdots+n_s > sk+1$. Then we have*

$$\begin{aligned}
2^{n_1+\cdots+n_s-2sk} q \sum_{l=0}^{n_1+\cdots+n_s-sk} & (-1)^l B_{l,n_1+\cdots+n_s-sk}(\frac{1}{2}) B_{l+sk}(q^{-1}) \\
&= \sum_{l=0}^{sk} B_{l,sk}(\frac{1}{2}) (-1)^{l+sk} B_{n_1+\cdots+n_s-l}(q).
\end{aligned}$$

3. Further remarks

In this ection, we assume that $q \in \mathbb{C}$ with $|q| < 1$. The q-Euler polynomials and q-Bernoulli polynomials are defined by the generating functions as follows:

$$F_q^E(t,x) = \frac{2}{qe^t+1} e^{xt} = \sum_{n=0}^{\infty} E_n(q|x) \frac{t^n}{n!}, \ |t + logq| < \pi,$$

and

$$F_q^B(t,x) = \frac{t}{qe^t-1} e^{xt} = \sum_{n=0}^{\infty} B_n(q|x) \frac{t^n}{n!}, \ |t + logq| < 2\pi. \qquad (38)$$

In the special case $x = 0$, $E(0|q) = E_n(q)$ are called the n-th q-Euler numbers and $B_n(0|q) = B_n(q)$ are also called the n-th q-Bernoulli numbers.
As usual convention, let us define $F_q^B(t, 0) = F_q^B(t)$ and $F_q^E(t, 0) = F_q^E(t)$.
For $s \in \mathbb{C}$, we have

$$\begin{aligned}\frac{q}{\Gamma(s)}\int_0^\infty F_q^B(-t,1)t^{s-2}dt &= \frac{q}{\Gamma(s)}\int_0^\infty \frac{e^{-t}}{1-qe^{-t}}t^{s-1}dt \\ &= \sum_{m=0}^{\infty} q^{m+1}\frac{1}{\Gamma(s)}\int_0^\infty e^{-(m+1)t}t^{s-1}dt \\ &= \sum_{m=0}^{\infty}\frac{q^{m+1}}{(m+1)^s} = \sum_{m=1}^{\infty}\frac{q^m}{m^s}.\end{aligned} \tag{39}$$

From (39), we define$q$-zeta function as follows:

$$\zeta_q(s) = \sum_{m=1}^{\infty}\frac{q^m}{m^s}, \text{ for } \Re(s) > 1.$$

Note that $\zeta_q(s)$ has meromorphic continuation to the whole complex s-plane with a simple pole at $s = 1$.
By (38), (39) and elementary complex integral, we get

$$\zeta_q(1-n) = (-1)^n\frac{qB_n(1|q)}{n} = \begin{cases} -(1+B_1(q)) & \text{if } \ n = 1, \\ (-1)^n\frac{B_n(q)}{n} & \text{if } \ n > 1. \end{cases} \tag{40}$$

From (38), we note that

$$\left(\frac{2}{qe^t+1}\right)\left(\frac{t}{qe^t-1}\right) = \frac{2t}{q^2e^{2t}-1} = \sum_{n=0}^{\infty}2^nB_n(q^2)\frac{t^n}{n!},$$

and

$$\left(\frac{2}{qe^t+1}\right)\left(\frac{t}{qe^t-1}\right) = e^{(B(q)+E(q))t} = \sum_{n=0}^{\infty}(B(q)+E(q))^n\frac{t^n}{n!}.$$

Thus, we have

$$2^nB_n(q^2) = (B(q)+E(q))^n = \sum_{l=0}^{n}\binom{n}{l}B_l(q)E_{n-l}(q), \tag{41}$$

with usual convention about replacing $B^n(q)$ by $B_n(q)$.
By (42), we get

$$B_n(q^2) = \frac{1}{2^n}\sum_{l=0}^{n}\binom{n}{l}B_l(q)E_{n-l}(q), (\text{ cf.}[16]). \tag{42}$$

From (40) and (41), we can derive the following equation:

$$\begin{aligned}\zeta_{q^2}(-n) &= (-1)^{n+1}\frac{q^2B_{n+1}(q^2)}{n+1} = -(-1)^n\frac{B_{n+1}(q^2)}{n+1} \\ &= -(-1)^n\frac{1}{2^{n+1}(n+1)}\sum_{l=0}^{n+1}\binom{n+1}{l}B_l(q^2)E_{n+1-l}(q^2).\end{aligned}$$

References

[1] M. Acikgoz, S. Araci, *A study on the integral of the product of several type Berstein polynomials*, IST Transaction of Applied Mathematics-Modelling and Simulation, 2010, Vol. 1, No.1(2), ISBN 1913-8342, 10–14.

[2] A. Bayad, T. Kim, B. Lee, S.-H. Rim, *Some identities on the Bernstein polynomials associated with q-Euler polynomials* (Communicated).

[3] S. Bernstein, *Démonstration du théorème de Weierstrass fondée sur le calcul des probabilités*, Commun. Soc. Math. Kharkow **13** (1912), 1–2.

[4] I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, *On the higher-order w-q-Genocchi numbers*, Adv. Stud. Contemp. Math. **19** (2009), 39–57.

[5] H. W. Gould, *A theorem concerning the Bernstein polynomials,* Mathematics Magazine, Vol. 31, No. 5 (1958), pp. 259264.

[6] N. K. Govil, V. Gupta, *Convergence of q-Meyer-König-Zeller-Durrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 97–108.

[7] L.C. Jang, W-J. Kim, Y. Simsek, *A study on the p-adic integral representation on $\mathbb{Z}_p$ associated with Bernstein and Bernoulli polynomials*, Advances in Difference Equations, Vol 2010 (2010), Article ID 163219, 6pp.

[8] K. I Joy, *Bernstein polynomials*, On-line Geometric Modelling Notes, http://en.wikipedia.org/wiki/Bernstein polynomails.

[9] T. Kim, *Barnes-type multiple q-zeta functions and q-Euler polynomials*, J. Physics A : Math. Theor., **43** (2010), 255201, 11pp.

[10] T. Kim, *An analogue of Bernoulli numbers and their congruences,* Rep. Fac. Sci. Engrg. Saga Univ. Math.22(1994), N.2, 21–26.

[11] T. Kim, *On the symmetries of the q-Bernoulli polynomials,* Abstract and Applied Analysis, Vol.2008(2008), Article ID 914367, 7pp.

[12] T. Kim, *A new approach to p-adic q-L-functions*, Advanced Studies in Contemporary Mathematics **12** (2006),N.1, 61–72.

[13] T. Kim, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients,* Russian Journal of Mathematical physics 15 (2008), 51–59.

[14] T. Kim, J. Choi, Y. H.Kim, *q-Bernstein polynomials associated with q-Stirling numbers and Carlit's q-Bernoulli numbers,* Abstract and Applied Analysis (in press).

[15] T. Kim, *Some identities on the q-Euler polynomials of higner order order and q-Stirling numbers by fermionic p-adic integral on $\mathbb{Z}_p$,* Russian Journal of Mathematical physics 16 (2009), 484–491.

[16] T. Kim, *Note on the Euler q-zeta functions*, Journal of Number Theory **129** (2009), 1798-1804.

[17] T. Kim, B. Lee, *Some identities of the Frobenius-Euler polynomials,* Abstract and Applied Analysis 2009 (2009), Article ID 639439,7 pp.

[18] T. Kim, J. Choi, Y.H. Kim, C. S. Ryoo, *On the fermionic p-adic integral representation of Bernstein polynomials associated with Euler numbers and polynomials,* Journal of inequalities and Applications, Vol 2010 (2010), Article ID 864249, 12 pp.

[19] B. A. Kupershmidt, *Reflection symmetries of q-Bernoulli polynomials,* J. Nonlinear Math. Phys. 12(2005), 412-422.

[20] G.G. Lorentz, *Bernstein polynomials,* (2nd edition) printed in USA (1986).

[21] S-H. Rim, J-H. Jin, E-J. Moon, S-J. Lee, *On multiple interpolation functions of the q-Genocchi polynomials,* Journal of Inequalities and Applications, Vol 2010 (2010), Article ID 351419, 13 pp..

[22] S. Zorlu, H. Aktuglu, and M. A. Ozarslan, *An estimation to the solution of an initial value problem via q-Bernstein polynomials,* Journal of Computational Analysis and Applications, Vol. 12, N. 3, (2010), 637645.

Abdelmejid Bayad. Département de mathématiques, Université d'Evry Val d'Essonne, Bd. F. Mitterrand, 91025 Evry Cedex, France,
E-mail address: `abayad@maths.univ-evry.fr`

Taekyun Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `tkkim@kw.ac.kr`